

We have blown this up so you can see how dangerous this stuff really is!
It is not safe for any living thing!



Ready Mixed Baitbits KILLS MICE AND RATS

Kills Warfarin-Resistant House Mice and Warfarin-Resistant Norway Rats.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: For control of House Mice, Norway Rats, and Roof Rats in and around homes, industrial, commercial, agricultural and public buildings. d-CON Ready Mixed Baitbits may also be used in and around transport vehicles (ships, trains, aircraft) and related port or terminal buildings. Do not use in sewers. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS: Place trays in or near points where you have seen mice or rat signs such as droppings, runways, burrows, or gnawing marks. Once areas requiring baiting have been identified, proceed as follows:

APPLICATION DIRECTIONS:

1. Place one ready-to-use bait tray in each potential feeding location. Place trays in dark, out-of-the-way locations, where rodents are likely to find them.

FOR HOUSE MICE: Use only one tray per location.

FOR NORWAY AND ROOF RATS: Start with one tray per location. Add trays, up to a maximum of four per location, if you determine that there is high rat activity at the location.

2. For best results, leave trays undisturbed for at least two days after placement. However, if contents have been scattered or mostly consumed sooner, replace trays. Clean up spilled bait. Check bait every two days, replacing trays as needed or until signs of rodent activity cease. Trays not fed from for five consecutive days may be relocated as needed. When there are no longer any signs of rodent activity, dispose of trays properly.

For complete control, continue baiting for at least 15 days. If you are in an area where there is a danger of reinfestation from adjoining property, permanent bait stations should be maintained. Stations should be checked periodically.

PRECAUTIONARY STATEMENTS:

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful or fatal if swallowed. Keep away from humans, domestic animals, and pets. Wash hands after handling bait. **IF BAIT IS EATEN BY HUMANS, CALL A PHYSICIAN AT ONCE.** For 24 hour emergency assistance, call your local Poison Control Center. **IF BAIT IS EATEN BY ANIMALS OR PETS:** Call your local veterinarian.

NOTE TO PHYSICIAN AND VETERINARIAN: This product may reduce the clotting ability of the blood and cause hemorrhaging. The anticoagulant action of this product may produce prolonged prothrombin times for 20 to 30 days after exposure. If poisoning occurs, intramuscular and oral administration of Vitamin K₁ are indicated, as in poisoning from overdose of dicumarol (bishydroxy coumarin). **FOR HUMAN CASES:** Vitamin K₁ is antidotal at doses of 10 to 20 mg (not mg/kg). Repeated treatments may need to be given for up to 30 days (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary. **FOR ANIMAL CASES:** Vitamin K₁ is antidotal at 5 mg/kg intramuscularly. Oral Vitamin K₁ should be given for up to 30 days at 5 mg/kg (based on monitoring of prothrombin times). In severe cases, blood transfusions may be necessary.

ENVIRONMENTAL HAZARDS:

This product is toxic to fish, birds and wildlife. This product can pose a secondary hazard to birds of prey and mammals. Do not apply directly to water.

STORAGE AND DISPOSAL:

STORAGE: Store in original container in a dry place inaccessible to children and pets.

DISPOSAL: Do not reuse empty container. Securely wrap contained and any unused bait in newspaper and discard in trash.

